

Journal of Oceanography

Volume 49 (1993)

Contents

Number 1

| | |
|---|--------|
| O. Katoh: Detailed Current Structures over the Continental Shelf off the San'in Coast in Summer | 1-16 |
| O. Katoh: Detailed Current Structures in the Eastern Channel of the Tsushima Strait in Summer | 17-30 |
| S. Tsunogai, M. Tsuzuki, S. Watanabe and N. Yoshida: Radiocarbon Anomaly Found in Aquicultural Scallops Suspended in Coastal Sea | 31-37 |
| T. Egawa, Y. Nagata and S. Sato: Seasonal Variation of the Current in the Tsushima Strait Deduced from ADCP Data of Ship-of-Opportunity | 39-50 |
| M. Nishimura, K. Kita-Tsukamoto, K. Kogure, K. Ohwada and U. Simidu: A New Method to Detect Viable Bacteria in Natural Seawater Using 16SrRNA Oligonucleotide Probe | 51-56 |
| H. Takeoka, O. Matsuda and T. Yamamoto: Processes Causing the Chlorophyll <i>a</i> Maximum in the Tidal Front in Iyo-Nada, Japan | 57-70 |
| Y. Isoda and T. Murayama: Diurnal Shelf Waves off Hamada on San'in Coast | 71-88 |
| S.-K. Yang, Y. Nagata, K. Taira and M. Kawabe: Southward Intrusion of the Intermediate Oyashio Water along the East Coast of the Boso Peninsula. I. Coastal Salinity-Minimum-Layer Water off the Boso Peninsula | 89-114 |

Short Contribution

| | |
|---|---------|
| S. Ohtsuka, N. Kubo, M. Okada and K. Gushima: Attachment and Feeding of Pelagic Copepods on Larvacean Houses | 115-120 |
| J. Yoshida, H. Sudo, M. Matsuyama, Y. Kurita and Y. Mine: Japan-Equator XBT Sections in Late November 1989 and in Early December 1991 | 121-129 |

Number 2

| | |
|--|---------|
| S.-C. Wen, P.-F. Guo and D.-C. Zhang: Analytically Derived Wind-Wave Directional Spectrum. Part 1. Derivation of the Spectrum | 131-147 |
| S.-C. Wen, P.-F. Guo, D.-C. Zhang, C.-L. Guan and H.-G. Zhan: Analytically Derived Wind-Wave Directional Spectrum. Part 2. Characteristics, Comparison and Verification of Spectrum | 149-172 |
| S.-K. Yang, Y. Nagata, K. Taira and M. Kawabe: Southward Intrusion of the Intermediate Oyashio Water along the East Coast of the Boso Peninsula, Japan. II. Intrusion Events into Sagami Bay | 173-191 |
| T. Y. Tang and Y. J. Yang: Low Frequency Current Variability on the Shelf Break Northeast of Taiwan | 193-210 |

| | |
|--|---------|
| T. Fujita, T. Inada and Y. Ishito: Density, Biomass and Community Structure of Demersal Fishes off the Pacific Coast of Northeastern Japan | 211-229 |
| A. Maeda, T. Yamashiro and M. Sakurai: Fluctuation in Volume Transport Distribution Accompanied by the Kuroshio Front Migration in the Tokara Strait | 231-245 |

Number 3

Special Section: The International Symposium on Global Change (IGBP)

| | |
|--|---------|
| S. Tsunogai: Introduction | 247 |
| T. Yanagi, T. Saino, T. Ishimaru and S. Uye: A Carbon Budget in Tokyo Bay | 249-256 |
| C.-T. A. Chen: Anthropogenic CO ₂ Distribution in the North Pacific Ocean | 257-270 |
| K. Kawamura and K. Usukura: Distributions of Low Molecular Weight Dicarboxylic Acids in the North Pacific Aerosol Samples | 271-283 |
| S. Montani and Y. Mishima: The Role of Nuta (Large Amorphous Particles) as a Nutrient Regenerator in Osaka Bay | 285-293 |
| E. Ohtaki, E. Yamashita and F. Fujiwara: Carbon Dioxide in Surface Seawaters of the Seto Inland Sea, Japan | 295-303 |
| S. Tsunogai, T. Ono and S. Watanabe: Increase in Total Carbonate in the Western North Pacific Water and a Hypothesis on the Missing Sink of Anthropogenic Carbon | 305-315 |

Regular Section

| | |
|--|---------|
| T. Matsuura and M. Kamachi: The Evolution of Isolated Vortices Interacted with Steep Slope | 317-352 |
| A. Maruyama, N. Mita and T. Higashihara: Particulate Materials and Microbial Assemblages around the Izena Black Smoking Vent in the Okinawa Trough | 353-367 |

Number 4

| | |
|--|---------|
| H. Takeoka, H. Akiyama and T. Kikuchi: The Kyucho in the Bungo Channel, Japan—Periodic Intrusion of Oceanic Warm Water | 369-382 |
| A. Kaneko, N. Gohda, W. Koterayama, M. Nakamura, S. Mizuno and H. Furukawa: Towed ADCP Fish with Depth and Roll Controllable Wings and Its Application to the Kuroshio Observation | 383-395 |
| K. Taira, S. Kitagawa, H. Otake and T. Asai: Observation of Temperature and Velocity from a Surface Buoy Moored in the Shikoku Basin (OMLET-88)—An Oceanic Response to a Typhoon | 397-406 |
| I. Yuasa, E. Hashimoto and H. Ueshima: Nitrogen and Phosphorus Distributions across the Thermohaline Front in Kii Channel in Winter | 407-424 |
| N. Takematsu, Y. Sato, Y. Kato and S. Okabe: Factors Regulating the Distribution of Elements in Marine Sediments Predicted by a Simulation Model | 425-441 |
| Y. Isoda and S. Saitoh: The Northward Intruding Eddy along the East Coast of Korea | 443-458 |
| K. Furuya, K. Takahashi and H. Iizumi: Wind-Dependent Formation of Phytoplankton Spring Bloom in Otsuchi Bay, a Ria in Sanriku, Japan | 459-475 |

Number 5

| | |
|--|---------|
| P. Chang: A Note on Solitary Waves along Potential Vorticity Fronts on an f -Plane | 477-489 |
| H. Takeoka and H. Murao: Tidal Currents Influenced by Topographic Eddies in Uchiumi Bay | 491-501 |
| T. Yanagi and S. Takahashi: Seasonal Variation of Circulations in the East China Sea and the Yellow Sea | 503-520 |
| A. Yoshimori: Rossby Number Dependence on Meander Dynamics of a Potential Vorticity Front | 521-533 |
| Y. Isoda: Generation Mechanism of Higher Mode Nondispersive Shelf Waves by Wind Forcing | 535-549 |
| P. Thadathil, A. Shikauchi, Y. Sugimori and M. Kubota: A Statistical Method to Get Surface Level Air-Temperature from Satellite Observations of Precipitable Water | 551-558 |
| E. Yamashita, F. Fujiwara, X. Liu and E. Ohtaki: Measurements of Carbon Dioxide in the Seto Inland Sea of Japan | 559-569 |
| R. Suzuki, M. Takahashi, K. Furuya and T. Ishimaru: Simplified Technique for the Rapid Determination of Phytoplankton Pigments by Reverse-Phase High-Performance Liquid Chromatography | 571-580 |
| W.-S. Chuang, H.-W. Li, T. Y. Tang and C.-K. Wu: Observations of the Countercurrent on the Inshore Side of the Kuroshio Northeast of Taiwan | 581-592 |
| Y. Sekine: Topographic Effect of a Marine Ridge on the Spin-Down of a Cyclonic Eddy with Mean Flow | 593-606 |

Number 6

| | |
|--|---------|
| A. Kasai, S. Kimura and T. Sugimoto: Warm Water Intrusion from the Kuroshio into the Coastal Areas South of Japan | 607-624 |
| E. Tanoue: Distributional Characteristics of DOC in the Central Equatorial Pacific | 625-639 |
| Y. Koike, Y. Nakaguchi, K. Hiraki, T. Takeuchi, T. Kokubo and T. Ishimaru: Species and Concentrations of Selenium and Nutrients in Tanabe Bay during Red Tide due to <i>Gymnodinium nagasakiense</i> | 641-656 |
| T. Naganuma and H. Seki: Abundance and Productivity of Bacterioplankton in a Eutrophication Gradient of Shimoda Bay | 657-665 |
| H. Akiyama and S. Saitoh: The <i>Kyuchio</i> in Sukumo Bay Induced by Kuroshio Warm Filament Intrusion | 667-682 |
| M. Matsuyama, S. Ohta, T. Hibiya and H. Yamada: Strong Tidal Currents Observed near the Bottom in the Suruga Trough, Central Japan | 683-696 |
| M. Ikeda: Sequential Updating Assimilation of Simulated Geosat Altimeter Data from Mesoscale Features into a Two-Layer Quasi-Geostrophic Model | 697-718 |

